

The Beta Cephei instability domain for the new solar composition and with new OP opacities

A. A. Pamyatnykh,^{1,2,3} W. Ziomek⁴¹ Copernicus Astronomical Center, Bartycka 18, 00-716 Warsaw, Poland² Institute of Astronomy, Pyatnitskaya Str. 48, 109017 Moscow, Russia³ Institut für Astronomie, Türkenschanzstrasse 17, 1180 Vienna, Austria⁴ Astronomical Institute, Kopernika 11, 51-622 Wrocław, Poland

The recent revision of the solar chemical composition (A04: Asplund et al. 2005) leads to a decrease of about 40% in the C, N, O, Ne abundances and to a $\sim 20\%$ decrease of Fe and some other metal abundances in comparison with older abundances (GN93: Grevesse & Noels 1993), as shown in Fig. 1.




Figure 1: The new solar abundances in comparison with the older ones.

We tested the effects of these modifications of the heavy element abundances on the instability of β Cephei models. For opacities, the newest data from the Opacity Project (Seaton 2005) were used. Fig. 2 shows that the β Cephei instability domain in the HRD, when computed with new data for $Z = 0.012$ (revised solar value), is very similar to the instability domain computed with the OPAL opacities (Iglesias & Rogers 1996) for older solar metallicities and $Z = 0.02$. For the older data and assuming $Z = 0.012$, we obtain only weak β Cep instability (Pamyatnykh 1999). Two effects are responsible for stronger instability when using the new data: (i) The metal opacity bump in the OP case is located slightly deeper in the star than that in the OPAL case, which results in more effective driving; (ii) at a fixed Z value, the new Fe-group abundances are higher than the older ones because the Z value is determined mainly by the abundances of C, N, O, and Ne (see Fig. 1).




Figure 2: The new β Cephei instability domain in the main-sequence band (OP opacity, A04 mixture, $Z = 0.012$) compared with the older one (OPAL GN93, $Z = 0.02$, see Pamyatnykh 1999). 29 bright variables from Stankov & Handler (2005) with $m_V < 6.0$ and well-measured Hipparcos parallaxes are plotted.

Acknowledgments. AAP acknowledges partial financial support from the HELAS project and from the Polish MNiI grant No. 1 P03D 021 28.

References

- Asplund M., Grevesse N., Sauval A. J., 2005, in Barnes III T. G., Bash F. N., eds, ASP Conf. Ser. Vol. 336, The Solar Chemical Composition. Astron. Soc. Pac., San Francisco, p. 25
- Grevesse N., Noels A., 1993, in Pratz N., Vangioni-Flam E., Casse M., eds, Origin and Evolution of the Elements. Cambridge Univ. Press, Cambridge, p. 15
- Iglesias C. A., Rogers F. J., 1996, ApJ, 464, 943
- Pamyatnykh A. A., 1999, Acta Astron., 49, 119
- Seaton M. J., 2005, MNRAS, 362, L1
- Stankov A., Handler G., 2005, ApJS, 158, 193